

Short communication

Isolation of a new polymorphic TC maize microsatellite located on the long arm of chromosome 10

Nurit Katzir^{1,*}, Yael Danin-Poleg¹, Yaakov Tadmor¹, Avri Bar-Zur¹ and Perry B. Cregan²

¹Department of Vegetable Crops, Agricultural Research Organization, Northern Research Center, Newe Ya'ar, P.O.B. 90000 Haifa 31900, Israel (*author for correspondence); ²USDA-ARS, BAZRC-West, Beltsville, MD 20705, USA

Received 14 November 1995; accepted in revised form 24 April 1996

Abstract

Simple sequence repeats (SSR), also called microsatellites, were previously proved to be an important class of DNA markers. The isolation, mapping and designing of primers to the flanking regions of a new maize SSR is reported. The new marker, with a core motif of (TC)₁₂, designated as MZETC34, was mapped to the long arm of chromosome 10.

Microsatellites, also called simple sequence repeats (SSR), have been described as 'second generation', polymorphic DNA markers [5]. They have been recommended as the standard markers used to prepare highly saturated genetic maps of any eukaryote genome [2]. A map based on SSR consists of a list of oligonucleotide sequences. Once published, an SSR becomes available to a wide range of potential users, and does not require shipping or storing of cloned probes. Akkaya et al. [1] assessed SSR length polymorphism in the soybean (*Glycine max* (L.) Merr.) and, for the first time, proved the usefulness of this marker in a higher plant. The presence, informativeness and wide application of SSRs have, since, been proven in other plant species including rice [12, 13], *Brassica napus* [7], barley [9], sunflower [4], *Arabidopsis thaliana* [3] and wheat [6]. The first six SSRs in maize were described and mapped by Senior and Heun [10].

Here, we report the isolation, mapping and designing of primers to the flanking regions of a new maize microsatellite, MZETC34, with a core motif (TC)₁₂. The microsatellite was isolated from a genomic library prepared using DNA from the inbred line J105. *Sau3AI* fragments, 200–400 bp long, were cloned in the pBluescript II KS+ plasmid vector (Stratagene, La Jolla, CA). The library was screened as described by Akkaya et al. [1] using (CT)₈ probe, and a TC positive clone designated MZETC34 was selected and

sequenced. Unique primers designed to the flanking regions of the (TC)₁₂ core element (Figure 1) were used to amplify corn DNA by the polymerase chain reaction (PCR), producing an amplification product which was 104 bp long in the J105 inbred line. Nine length alleles were detected when these primers were screened on 19 maize genotypes and analyzed on a sequencing gel. The inbred lines IL731A and W6786 [11] were among the polymorphic genotypes. The amplification product of IL731A was 100 bp long while that of W6786 was 104 bp long (Figure 1).

Seventy seven F_{2:3} families of the cross IL731a × W6786 were scored for MZETC34 segregation (Figure 1). Twenty one were homozygotes carrying the 100 bp allele, 34 were heterozygotes, and 22 were homozygotes carrying the 104 bp allele. The results fitted the 1:2:1 ratio of codominant segregation with a χ^2 value of 1.08 ($P = 0.583$). The sweet corn linkage map was recently established using this population [11]. In order to localize MZETC34 on the sweet corn map, the linkage relationships between MZETC34 and 91 RFLP markers were analyzed with MAPMAKER [8] using a LOD score of 3.0 and a recombination fraction of 0.4. Employing this procedure MZETC34 was located on the long arm of chromosome 10, situated between the RFLP markers umc 152 and npi 582 (Figure 2).




Figure 1. Radiolabelled PCR products amplified in the F₂ progeny of the cross IL731A × W6876. P₁: IL731A; P₂: W6876; lanes 1–35: F₂ plants. The sequencing ladder is M13mp18 ssDNA (GATC). Amplification primers flanking the (TC)₁₂ motif: CCAAAACACAAGTCTGAACTC (forward); CTGTGAAAGGGATGCAAG (reverse).




Figure 2. A linkage map of RFLP and the MZETC34 SSR on chromosome 10.

References

1. Akkaya MS, Bhagwat AA, Cregan PB: Length polymorphism of simple sequence repeat DNA in soybean. *Genetics* 132: 1131–1139 (1992).

2. Beckmann JS, Soller M: Toward a unified approach to genetic mapping of eukaryotes based on sequence tagged microsatellite sites. *Bio/technology* 8: 30–32 (1990).

3. Bell CJ, Ecker JR: Assignment of 30 microsatellite loci to the linkage map of *Arabidopsis*. *Genomics* 19: 137–144 (1994).

4. Brunel D: A microsatellite marker in *Helianthus annuus* L. *Plant Mol Biol* 24: 397–400 (1994).

5. Davies K: Of mice and men (and cows and cats). *Nature* 361: 478 (1993).

6. Devos KM, Bryan GJ, Collins AJ, Stephenson P, Gale MD: Application of two microsatellite sequences in wheat storage proteins as molecular markers. *Theor Appl Genet* 90: 247–252 (1995).

7. Lagercrantz U, Ellegren H, Andersson L: The abundance of various polymorphic microsatellite motifs differs between plants and vertebrates. *Nucl Acids Res* 21: 1111–1115 (1993).

8. Lander ES, Green P, Abrahamson J, Barlow A, Daly MJ, Lincoln SE, Newburg L: MAPMAKER: an interactive computer package for constructing primary genetic linkage maps of experimental and natural populations. *Genomics* 1: 174–181 (1987).

9. Saghai-Maroo MA, Biyashev RM, Yang GP, Zhang Q, Allard RW: Extraordinarily polymorphic microsatellite DNA in barley: species diversity, chromosomal locations, and population dynamics. *Proc Natl Acad Sci USA* 91: 5466–5470 (1994).

10. Senior ML, Heun M: Mapping maize microsatellites and polymerase chain reaction confirmation of the targeted repeats using a CT primer. *Genome* 36: 884–889 (1993).

11. Tadmor Y, Azanza F, Han T, Rocheford TR, Juvik JA: RFLP mapping of the *sugary enhancer 1* gene in maize. *Theor Appl Genet* 91: 489–494 (1995).

12. Wu KS, Tanksley SD: Abundance, polymorphism and genetic mapping of microsatellites in rice. *Mol Gen Genet* 241: 225–235 (1993).

13. Zhao X, Kochert G: Phylogenetic distribution and genetic mapping of a (GGC)_n microsatellite from rice (*Oryza sativa* L.). *Plant Mol Biol* 21: 607–614 (1993).